

Second order transport from anomalies

Sayantani Bhattacharyya,^a Justin R. David^b and Somyadip Thakur^b.

^a *Physics Department,
Ramkrishna Mission Vivekananda University,
Belur Math, Howrah 711202, India.*

^b *Centre for High Energy Physics, Indian Institute of Science,
C.V. Raman Avenue, Bangalore 560012, India.*

sayanta@rkmvu.ac.in, justin,somyadip@cts.iisc.ernet.in

ABSTRACT: We study parity odd transport at second order in derivative expansion for a non-conformal charged fluid. We see that there are 27 parity odd transport coefficients, of which 12 are non-vanishing in equilibrium. We use the equilibrium partition function method to express 7 of these in terms of the anomaly, shear viscosity, charge diffusivity and thermodynamic functions. The remaining 5 are constrained by 3 relations which also involve the anomaly. We derive Kubo formulae for 2 of the transport coefficients and show these agree with that derived from the equilibrium partition function.

Contents

1. Introduction	1
1.1 Summary of the results	3
2. Anomalous transport from equilibrium partition function	7
2.1 The equilibrium partition function method	7
2.2 Partition function analysis: parity-odd sector at second order	9
3. Kubo formula for the transport coefficients Φ_1, Φ_2	18
4. Conclusions	22

1. Introduction

Fluid dynamics is an effective description of near-equilibrium systems. Properties of a fluid system are always slowly varying compared to some intrinsic length scale, for example a mean free path. This length scale is determined by the details of the underlying microscopic theory. The fundamental variables in fluid dynamics are local velocities $u^\mu(x)$, temperature $T(x)$, and all the other conserved charges or their chemical potentials $\mu^a(x)$. The conservation equations for the stress tensor $T^{\mu\nu}$ and the other conserved currents $J^{a\mu}$ govern the time evolution in fluid dynamics.

The stress tensor and the conserved currents are related to the fluid variables $\{u^\mu, T, \mu^a\}$ via constitutive relations. Since fluid systems are always slowly varying it is appropriate to organize the constitutive relations in terms of a derivative expansion of the fluid variables. At every order in the derivative expansion, the independent terms of the constitutive relation are constructed out of the independent derivatives of the fluid variables. The independent terms in the constitutive relation are multiplied by coefficients which are functions of temperature and chemical potentials. These coefficients are called transport coefficients. In this paper we will study the transport coefficients that occur in the parity odd sector, at second order in the derivative expansion. These are terms constructed out of various derivatives in the fluid variables which are odd under parity. We will consider relativistic fluid systems with one additional conserved current.

It is usually difficult to compute transport coefficients from the microscopic theory and they are generically determined from experiments. However parity odd

transport coefficients which occur at the first order in the derivative expansion have been related to quantum anomalies of the microscopic theory [1, 2, 3, 4, 5, 6, 7]¹. Our goal is to see if a similar phenomenon occurs at the second order in the derivative expansion of the constitutive relations. One motivation to consider second order fluid dynamics is that first order fluid dynamics is known to have problems with causality and numerical stability. Transport coefficients which occur at second order provide important constraints for spectral densities through sum rules [9]. Parity odd transport coefficients at second order affect the dispersion relation of chiral modes [10]². This phenomenon has important experimental consequences like the spatial separation of particles of different chirality. In the relativistic context this phenomenon was first observed holographically [11] and then understood due to the presence of a parity odd transport coefficient at second order by [12].

Parity odd transport coefficients at second order has been studied earlier in [12] for conformal fluids. They used the principle that parity odd terms which are even under time-reversal invariance should not contribute to local entropy production. With this principle they could constrain these transport coefficients and determine some of them. We will use the method developed in [13, 14] to determine and constrain the parity odd transport coefficients. We consider non-conformal fluids in $3 + 1$ dimensions which admits one anomalous charge current. This method is based on the requirement that the fluid equations have to be consistent with any equilibrium partition function. Therefore the approach first relies on the physical requirement of the existence of equilibrium. More precisely:

- In a time independent background, that is a space-time metric with a time like Killing vector and background gauge fields independent of the time direction, any fluid equation will admit a time independent solution.

The second assumption is:

- The stress tensor and the charge current evaluated on this time independent solution can be obtained from the partition function by varying it with respect to the background metric and the gauge field.

This method is implemented as follows.

1. We first classify all the parity odd transport coefficients till the second order in the derivative expansion of the stress tensor and the charge current using symmetries.
2. We then evaluate the stress tensor and the charge current on the equilibrium fluid configuration to the second order in the derivative expansion.

¹See [8] for a recent review with a complete list of references.

²We thank Yashodhan Hatwalne for bringing this reference to our attention.

3. The equilibrium partition function is written to the second order in derivative expansion taking all the parity odd terms into consideration. This is also done based on symmetries
4. This stress tensor and the charge current obtained from the equilibrium partition function is required to agree with that obtained from the stress tensor evaluated on the equilibrium fluid configuration.

From the description of the method it is clear that only transport coefficients which do not vanish in the equilibrium fluid configuration will be constrained or determined. We will see that totally there are 27 parity odd transport coefficients. Out of these 12 do not vanish in the equilibrium fluid configuration. Among the 12, we determine 7 which we label as $\Phi_i, i = 1, \dots, 6$ and Δ_2 . and show that they are related to the anomaly. The rest of the 5 are constrained by 3 relations. These relations are also involve the anomaly. The results are summarized in (1.9) and (1.10). We will then use the Kubo formula to derive two of the transport coefficients Φ_1, Φ_2 and show that it agrees with the equilibrium partition function method. The remaining transport coefficients seem to be related to three point functions.

The organization of the paper is as follows. In the rest of the introduction we summarize our main results. In section 2 we implement the method of [13] to relate the transport coefficients which do not vanish in equilibrium to that of the anomaly. In section 3 we use the Kubo formula to derive two of the transport coefficients.

1.1 Summary of the results

As we mentioned earlier, the aim of this note is to constrain the parity odd second order transport coefficients of an anomalous charged fluid in the presence of background electric and magnetic fields. To define the transport coefficients unambiguously we first must have an unambiguous definition of fluid variables that is the velocities, the temperature and the chemical potentials. $\{u^\mu, T, \mu\}$ We will work in Landau frame which is defined by the following two conditions for the charged fluid.

$$J^\mu u_\mu = -q, \quad T^{\mu\nu} u_\mu = -E u^\nu. \quad (1.1)$$

Let us now consider the expansion of the charge current J^μ and the stress tensor $T^{\mu\nu}$ in terms of the number of space time derivatives. This is given by

$$J^\mu = J_{(0)}^\mu + J_{(1)}^\mu + \dots, \quad T^{\mu\nu} = T_{(0)}^{\mu\nu} + T_{(1)}^{\mu\nu} + \dots, \quad (1.2)$$

where the subscript (i) refer to the number of space-time derivatives. The terms $J_{(i)}^\mu$ and $T_{(i)}^{\mu\nu}$ for $i \neq 0$ are all perpendicular to the velocity u^μ . The equations of motion for the fluid in presence of external electromagnetic field are given by the following

conservation laws

$$\begin{aligned}
\nabla_\mu T^{\mu\nu} &= \mathcal{F}^{\nu\mu} J_\mu + \frac{c_m}{2} \epsilon^{\alpha\beta\gamma\delta} \mathcal{F}_{\alpha\beta} R_{\gamma\delta}^{\mu\nu}, \\
\nabla_\mu J^\mu &= -\frac{C}{8} \epsilon^{\alpha\beta\gamma\delta} \mathcal{F}_{\alpha\beta} \mathcal{F}_{\gamma\delta} + \frac{c_m}{4} \epsilon^{\alpha\beta\gamma\delta} R_{\nu\alpha\beta}^\mu R_{\mu\gamma\delta}^\nu \\
&= C E_\mu B^\mu + \frac{c_m}{4} \epsilon^{\alpha\beta\gamma\delta} R_{\nu\alpha\beta}^\mu R_{\mu\gamma\delta}^\nu.
\end{aligned} \tag{1.3}$$

Here $E_\mu = \mathcal{F}_{\mu\nu} u^\nu$, $B^\mu = \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} u_\nu \mathcal{F}_{\alpha\beta}$ and C is the gauge anomaly coefficient and c_m is the coefficient of the mixed gauge-gravitational anomaly. Note that the terms proportional to the mixed anomaly are fourth order in derivatives, therefore they do not affect the analysis of the equations of motion to 2nd order in derivatives. However the gravitational anomaly does enter the discussion of the equilibrium partition function at the first order in the derivative expansion [15].

We will now state the known results for the form of the stress tensor and the current up to first order in the derivative expansion. For $i = 0$, the part with no space-time derivative the stress tensor and current is completely determined by thermodynamics. At first order in derivative expansion, that is $i = 1$, the form of the current and stress tensor is explicitly known [13], [1]. This form is consistent with all physical requirements in presence of anomaly and as well as an external electromagnetic field [13], [1]. The final result for the stress tensor and the charge current to first order in derivatives is given by

$$\begin{aligned}
[T_{(0)}]_{\mu\nu} &= (E + P) u_\mu u_\nu + P G_{\mu\nu}, \\
[T_{(1)}]_{\mu\nu} &= -2\eta \sigma^{\mu\nu} - \zeta \Theta P^{\mu\nu}, \\
J_{(0)}^\mu &= q u^\mu, \\
J_{(1)}^\mu &= \Delta V^\mu + \xi_l l^\mu + \xi_B B^\mu.
\end{aligned} \tag{1.4}$$

Here E is the energy density, P the pressure and q , the charge density and $G_{\mu\nu}$ is the background metric. The variables Θ , $\sigma^{\mu\nu}$, V^μ , l^μ and B^μ are all on-shell independent terms which are first order in derivatives. These are defined in table 1. The variables η , ζ , Δ , ξ_l and ξ_B refer to the first order transport coefficients.

Scalars (1)	Vectors (3)	Pseudo Vectors (2)	Tensors (1)
$\Theta = \nabla_\mu u^\mu$	$u^\mu \nabla_\mu u_\nu$ $P^{\mu\nu} \nabla_\nu \nu$ $V^\mu = (E^\mu - T P^{\mu\nu} \nabla_\nu \nu)$	$l^\mu = \epsilon^{\mu\nu\alpha\beta} u_\nu \partial_\alpha u_\beta$ $B^\mu = \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} u_\nu \mathcal{F}_{\alpha\beta}$	$\sigma_{\mu\nu} = \nabla_{\langle\mu} u_{\nu\rangle}$

Table 1: Data at 1st order in derivative

Throughout this paper, the symbol $A_{\langle\mu\nu\rangle}$ on any tensor $A_{\mu\nu}$ denotes the projected, traceless, symmetric part of the tensor.

$$A_{\langle\mu\nu\rangle} = P_{\mu}^{\alpha} P_{\nu}^{\beta} \left(\frac{A_{\alpha\beta} + A_{\beta\alpha}}{2} - \frac{P^{\gamma\theta} A_{\gamma\theta}}{3} G_{\alpha\beta} \right), \quad (1.5)$$

and $P^{\mu\nu}$ is the projector

$$P^{\mu\nu} = u^{\mu} u^{\nu} + G^{\mu\nu}. \quad (1.6)$$

The variable ν is related to the chemical potential by

$$\nu = \frac{\mu}{T}. \quad (1.7)$$

Let us now proceed to the stress tensor and the charge current at second order in derivatives which we denote as $[T_{(2)}]_{\mu\nu}$ and $J_{(2)}^{\mu}$. Purely from symmetry considerations, the number of independent transport coefficients upto second order is equal to total number of possible scalars which appear in the trace of the stress tensor, together with the number of possible vectors which appear in the current and the possible symmetric traceless tensors which appear in the traceless part of the stress tensor. But not all of them are independent, they can be related using the equations of motion given in (1.3). In Table 2 we have listed all the parity odd and on-shell independent scalars, vectors and tensors containing two space-time derivatives.

Pseudo-scalars (6)	Pseudo Vectors (9)	Pseudo-tensors (12)
$\mathcal{S}_1 = l^{\mu} \partial_{\mu} \nu$ $\mathcal{S}_2 = B^{\mu} \partial_{\mu} \nu$ $\mathcal{S}_3 = l^{\mu} \partial_{\mu} T$ $\mathcal{S}_4 = B^{\mu} \partial_{\mu} T$ $\mathcal{S}_5 = l^{\mu} V_{\mu}$ $\mathcal{S}_6 = B^{\mu} V_{\mu}$	$\mathcal{V}_{(1)}^{\mu} = \epsilon^{\mu\nu\alpha\beta} u_{\nu} B_{\alpha} l_{\beta}$ $\mathcal{V}_{(2)}^{\mu} = \epsilon^{\mu\nu\alpha\beta} u_{\nu} (\partial_{\alpha} \nu) (\partial_{\beta} T)$ $\mathcal{V}_{(3)}^{\mu} = \Theta l^{\mu}$ $\mathcal{V}_{(4)}^{\mu} = (u \cdot \nabla) B^{\mu}$ $\mathcal{V}_{(5)}^{\mu} = \sigma_{\nu}^{\mu} l^{\nu}$ $\mathcal{V}_{(6)}^{\mu} = \Theta B^{\mu}$ $\mathcal{V}_{(7)}^{\mu} = \sigma_{\nu}^{\mu} B^{\nu}$ $\mathcal{V}_{(8)}^{\mu} = \epsilon^{\mu\nu\alpha\beta} u_{\nu} (\partial_{\alpha} T) V_{\beta}$ $\mathcal{V}_{(9)}^{\mu} = \epsilon^{\mu\nu\alpha\beta} u_{\nu} (\partial_{\alpha} \nu) V_{\beta}$	$\tau_{\mu\nu}^{(1)} = \nabla_{\langle\mu} l_{\nu\rangle}$ $\tau_{\mu\nu}^{(2)} = \nabla_{\langle\mu} B_{\nu\rangle}$ $\tau_{\mu\nu}^{(3)} = l_{\langle\mu} \partial_{\nu\rangle} \nu$ $\tau_{\mu\nu}^{(4)} = B_{\langle\mu} \partial_{\nu\rangle} \nu$ $\tau_{\mu\nu}^{(5)} = l_{\langle\mu} \partial_{\nu\rangle} T$ $\tau_{\mu\nu}^{(6)} = B_{\langle\mu} \partial_{\nu\rangle} T$ $\tau_{\mu\nu}^{(7)} = l_{\langle\mu} V_{\nu\rangle}$ $\tau_{\mu\nu}^{(8)} = B_{\langle\mu} V_{\nu\rangle}$ $\tau_{\mu\nu}^{(9)} = u_{\theta} \sigma_{\langle\mu\alpha} (\partial_{\beta} T) \epsilon^{\theta\alpha\beta}{}_{\nu\rangle}$ $\tau_{\mu\nu}^{(10)} = u_{\theta} \sigma_{\langle\mu\alpha} (\partial_{\beta} \nu) \epsilon^{\theta\alpha\beta}{}_{\nu\rangle}$ $\tau_{\mu\nu}^{(11)} = u_{\theta} \sigma_{\langle\mu\alpha} V_{\beta} \epsilon^{\theta\alpha\beta}{}_{\nu\rangle}$ $\tau_{\mu\nu}^{(12)} = u_{\theta} \nabla_{\beta} \sigma_{\langle\mu\alpha} \epsilon^{\theta\alpha\beta}{}_{\nu\rangle}$

Table 2: Parity odd data at 2nd order in derivatives

From the table it can be seen that at second order in derivatives there are 27 parity odd transport coefficients which appear in the current and the stress tensor. Therefore the most general parity odd contributions at second order in stress tensor and current can be parametrized as follows.

$$\begin{aligned} [T_{(2)}]_{\mu\nu} &= \sum_{i=1}^{12} \Phi_i \tau_{\mu\nu}^{(i)} + P_{\mu\nu} \left[\sum_{i=1}^6 \chi_i \mathcal{S}_i \right], \\ J_{(2)}^\mu &= \sum_{i=1}^9 \Delta_i \mathcal{V}_{(i)}^\mu. \end{aligned} \quad (1.8)$$

Where $\tau_{\mu\nu}^{(i)}$, $\mathcal{S}_i$ and $\mathcal{V}_{(i)}^\mu$ are defined in Table 2. Our goal is to constrain the transport coefficients Φ_i , χ_i and Δ_i , using the existence of the equilibrium partition function. It can be shown among these 27 terms only 12 can be non-zero in a time independent equilibrium fluid configuration. They are the first 4 in the list of scalars $\mathcal{S}_1 \cdots \mathcal{S}_4$, the first 2 in the list of vectors $\mathcal{V}_{(1)}^\mu, \mathcal{V}_{(2)}^\mu$ and the first 6 in the list of tensors $\tau_{\mu\nu}^{(1)} \cdots \tau_{\mu\nu}^{(6)}$. Therefore the analysis using the equilibrium partition function can be used to constrain the 12 transport coefficients multiplying these non-vanishing terms. These are $\chi_1 \cdots \chi_4$ and Δ_1, Δ_2 and $\Phi_1 \cdots \Phi_6$. The final result of this analysis is the following.

$$\begin{aligned} \Phi_1 &= \eta b_1, \quad \Phi_2 = 2\eta b_2, \quad \Phi_3 = \eta \left(\frac{\partial b_1}{\partial \nu} \right), \quad \Phi_4 = 2\eta \left(\frac{\partial b_2}{\partial \nu} \right), \\ \Phi_5 &= \eta \left[-\frac{b_1}{T} + \frac{\partial b_1}{\partial T} \right], \quad \Phi_6 = 2\eta \left[-\frac{b_2}{T} + \frac{\partial b_2}{\partial T} \right], \end{aligned} \quad (1.9)$$

$$\begin{aligned} \Delta_2 &= -\frac{\Delta b_1}{2}, \\ T^2 R_1 \left[\chi_3 - \frac{\zeta}{2} \left(\frac{\partial b_1}{\partial T} - \frac{2b_1}{T} \right) \right] - R_2 \left[\chi_1 - \frac{\zeta}{2} \left(\frac{\partial b_1}{\partial \nu} - 2b_2 T \right) \right] &= 0, \\ T^2 R_1 \left[\chi_4 - \zeta \left(\frac{\partial b_2}{\partial T} - \frac{b_2}{T} \right) \right] + R_2 \left[\chi_2 - \zeta \left(\frac{\partial b_1}{\partial \nu} \right) \right] &= 0, \\ R_1 T \Delta_1 + \left[\chi_2 - \zeta \left(\frac{\partial b_2}{\partial \nu} \right) \right] - \frac{q}{(E+P)} \left[\chi_1 - \frac{\zeta}{2} \left(\frac{\partial b_1}{\partial \nu} - 2b_2 T \right) \right] &= 0, \end{aligned} \quad (1.10)$$

where

$$\begin{aligned} b_1 &= \frac{T^3}{E+P} \left(\frac{2C\nu^3}{3} - 4C_2\nu \right), \quad b_2 = \frac{T^2}{E+P} \left(\frac{C\nu^2}{2} - C_2 \right), \\ R_1 &= \left(\frac{\partial P}{\partial E} \right)_q, \quad R_2 = \left(\frac{\partial P}{\partial q} \right)_E. \end{aligned}$$

Note that C is the gauge anomaly coefficient and C_2 is related to the coefficient of the mixed gauge-gravitational anomaly in (1.3) by

$$C_2 = 8\pi^2 c_m. \quad (1.11)$$

Therefore we see that the coefficients $\Phi_1, \dots, \Phi_6$ and Δ_2 are determined in terms of the anomaly, the shear viscosity, the charge diffusivity Δ and the thermodynamic functions E, P, T, ν . The rest $\Delta_1, \chi_1, \dots, \chi_4$ are constrained by 3 relations which involve the anomaly.

2. Anomalous transport from equilibrium partition function

In sub-section 2.1 we will first we briefly outline the general procedure we use to relate parity odd transport at the second order in derivatives to the anomaly. This method has been described and analyzed in detail by [13] and [14]. It relies on the analysis of the equilibrium partition function. Thus we will be able to constrain only those transport coefficients which do not vanish in the equilibrium configuration. This section also will serve to introduce the notation and conventions used in the paper. In sub-section 2.2 we implement this method and derive the relations given in (1.9) and (1.10).

2.1 The equilibrium partition function method

We are interested in a fluid flow on a static background metric and a static external electromagnetic field. The most general static metric and gauge field can be written in the following form.

$$ds^2 = G_{\mu\nu} dx^\mu dx^\nu = -e^{2\sigma} (dt + a_i dx^i)^2 + g_{ij} dx^i dx^j, \quad (2.1)$$

Gauge Field : $\mathcal{A}_\mu dx^\mu$,

$\sigma, a_i, \mathcal{A}_0$ and $\mathcal{A}_\mu$ are all slowly varying functions of the spatial co-ordinates $(\vec{x})$. Our notations are as follows:

- Greek indices run from 1 to 4.
- Latin indices run from 1 to 3.
- All Greek indices are lowered or raised by the 4 dimensional metric $G_{\mu\nu}$ unless explicitly mentioned.
- All Latin indices are lowered or raised by the 3 dimensional metric g_{ij} unless explicitly mentioned.
- $\bar{\nabla}_\mu$ is covariant derivative with respect to the metric $G_{\mu\nu}$ and ∇_i is the covariant derivative with respect to the metric g_{ij} .
- For any tensor $A_{\mu\nu}$ the notation $A_{\langle\mu\nu\rangle}$ denotes the traceless symmetric part of the tensor, projected in direction perpendicular to the fluid velocity.

$$A_{\langle\mu\nu\rangle} = P_\mu^\alpha P_\nu^\beta \left(\frac{A_{\alpha\beta} + A_{\beta\alpha}}{2} - \frac{P^{\gamma\theta} A_{\gamma\theta}}{3} G_{\alpha\beta} \right),$$

where $P_{\mu\nu} \equiv u_\mu u_\nu + G_{\mu\nu}$ is referred to as the projector.

Our basic assumption is that in such a background any fluid equation will admit a time independent solution. The stress tensor and the current, evaluated on this time independent solution, can be generated by varying the partition function of the system with respect to the background metric and gauge field. If Z is the partition function at temperature T_0 then the stress tensor and the current evaluated on the equilibrium are given by the following formulae.

$$\begin{aligned}
T_{00}|_{\text{equilibrium}} &= -\frac{T_0 e^{2\sigma}}{\sqrt{-G}} \left[\frac{\delta Z}{\delta \sigma} \right] = \frac{T_0^2}{\sqrt{g}} \left[\frac{\delta Z}{\delta \bar{T}} \right], \\
T_0^i|_{\text{equilibrium}} &= \frac{T_0}{\sqrt{-G}} \left[\frac{\delta Z}{\delta a_i} - A_0 \frac{\delta Z}{\delta A_i} \right] = \frac{\bar{T}}{\sqrt{g}} \left[\frac{\delta Z}{\delta a_i} - A_0 \frac{\delta Z}{\delta A_i} \right], \\
T^{ij}|_{\text{equilibrium}} &= \frac{2T_0}{\sqrt{-G}} g^{il} g^{jm} \left[\frac{\delta Z}{\delta g^{lm}} \right] = \frac{2\bar{T}}{\sqrt{g}} g^{il} g^{jm} \left[\frac{\delta Z}{\delta g^{lm}} \right], \\
J_0|_{\text{equilibrium}} &= -\frac{T_0 e^{2\sigma}}{\sqrt{-G}} \left[\frac{\delta Z}{\delta A_0} \right] = -\frac{e^\sigma}{\sqrt{g}} \left[\frac{\delta Z}{\delta \bar{\nu}} \right], \\
J^i|_{\text{equilibrium}} &= \frac{T_0}{\sqrt{-G}} \left[\frac{\delta Z}{\delta A_i} \right] = \frac{\bar{T}}{\sqrt{g}} \left[\frac{\delta Z}{\delta A_i} \right],
\end{aligned} \tag{2.2}$$

where

$$\bar{T} = T_0 e^{-\sigma}, \quad \bar{\nu} = \frac{A_0}{T_0}, \quad A_i = \mathcal{A}_i - A_0 a_i.$$

The strategy we will adopt to constrain the parity odd coefficients which occur at the second order is the following:

1. Write down the most general partition function upto a given order in derivative expansion and consistent with all the symmetries. It will be a function of a_i , σ and $\mathcal{A}_i$ and their derivatives.
2. Vary the partition function Z , to obtain the most general possible expression for $T^{\mu\nu}|_{\text{equilibrium}}$ and $J^\mu|_{\text{equilibrium}}$.
3. Parametrize the most general possible fluid stress tensor and current up to some given order in derivative expansion using symmetries. This will give the maximum number of independent transport coefficients possible constrained only by symmetry. We have stated the results of this analysis already in equation (1.4) and (1.8).
4. Evaluate the most general fluid stress tensor and the current on the equilibrium solution. The final outcome will contain some of the unknown transport coefficients.
5. Equate the final outcome of the previous step with the stress tensor and current we have already obtained by varying the partition function.

6. This will express the transport coefficients which appear in the stress tensor and the current evaluated on equilibrium, in terms of the coefficients appearing in the partition function.
7. Eliminating the coefficients which appear in the partition function results in the desired relations among the transport coefficients.

One might wonder that to execute the fourth step, the equilibrium solution for the velocity, temperature and the other fluid variables needs to be independently found. But as it has been explained in [13] that using this method one can perturbatively determine both the solution and the transport coefficients in terms of the free functions appearing in the partition function.

2.2 Partition function analysis: parity-odd sector at second order

In this subsection we shall apply the general procedure described in 2.1 to the particular case of parity-odd sector of the charged fluid at second order in derivative expansion. The first contribution in parity-odd sector comes at first order in derivative expansion. This has been analyzed in detail in section 3 of [13]. We will not repeat the first order computation here but we will extensively use their result.

Stress tensor and current from the partition function

Since the transport coefficients we are interested in is parity-odd sector, we will restrict our attention to only the parity odd part of the partition function. The partition function $Z_{(2)}$ at second order in derivatives is a gauge invariant scalar functional of the background metric and gauge-fields.³ Hence we need to list all possible parity odd scalars that can be constructed from the metric functions and gauge fields that contain two space-derivatives. Note that since all the functions are time independent no time derivatives occur. There are four such scalars:

1. $\epsilon^{ijk} \partial_i \bar{\nu} f_{jk}$,
2. $\epsilon^{ijk} \partial_i \bar{T} f_{jk}$,
3. $\epsilon^{ijk} \partial_i \bar{\nu} F_{jk}$,
4. $\epsilon^{ijk} \partial_i \bar{T} F_{jk}$,

where $\bar{\nu} = \frac{A_0}{T_0}$, $\bar{T} = T_0 e^{-\sigma}$, $A_i = \mathcal{A}_i - A_0 a_i$ and $f_{jk} = \partial_j a_k - \partial_k a_j$, $F_{jk} = \partial_j A_k - \partial_k A_j$.

Therefore naively, the parity odd second order partition function at two derivatives can have 4 free parameters, but two of them can be related by total derivatives.

³Though the system is anomalous, all the effects of anomaly i.e. the anomalous transformation property of the partition function under the gauge transformation is accounted for in first order part $Z_{(1)}$. Therefore in $Z_{(2)}$ we need to consider only the gauge invariant scalars.

Ignoring the total derivative terms the most general second order partition function can be written as

$$Z_{(2)} = \int \sqrt{g} \left[M_1(\bar{T}, \bar{\nu}) \epsilon^{ijk} \partial_i \bar{\nu} F_{jk} + T_0 M_2(\bar{T}, \bar{\nu}) \epsilon^{ijk} \partial_i \bar{\nu} f_{jk} \right].$$

Varying this partition function with respect to the metric and the gauge fields and using (2.2) we get the following second order correction to the stress tensor and charge current.

$$\begin{aligned} [\Pi^{(2)}]_{00} &= T_0^2 \left[\left(\frac{\partial M_1}{\partial \bar{T}} \right) \epsilon^{ijk} \partial_i \bar{\nu} F_{jk} + T_0 \left(\frac{\partial M_2}{\partial \bar{T}} \right) \epsilon^{ijk} \partial_i \bar{\nu} f_{jk} \right], \\ [\Pi^{(2)}]_0^i &= 2T_0 \bar{T} \left(\frac{\partial M_2}{\partial \bar{T}} - \bar{\nu} \frac{\partial M_1}{\partial \bar{T}} \right) \epsilon^{ijk} (\partial_j \bar{T}) (\partial_k \bar{\nu}), \\ [\Pi^{(2)}]^{ij} &= 0, \end{aligned} \tag{2.3}$$

$$\begin{aligned} [j^{(2)}]_0 &= \frac{T_0}{\bar{T}} \left[\left(\frac{\partial M_1}{\partial \bar{T}} \right) \epsilon^{ijk} \partial_i \bar{T} F_{jk} + T_0 \left(\frac{\partial M_2}{\partial \bar{T}} \right) \epsilon^{ijk} \partial_i \bar{T} f_{jk} \right], \\ [j^{(2)}]^i &= 2\bar{T} \left(\frac{\partial M_1}{\partial \bar{T}} \right) \epsilon^{ijk} (\partial_j \bar{T}) (\partial_k \bar{\nu}). \end{aligned}$$

It is important to note the third equation in (2.3). The fact that the spatial components of the equilibrium stress tensor vanish at second order will serve as an important constraint in determining the transport coefficients.

Stress tensor from fluid dynamics

We have to evaluate the fluid dynamical stress tensor on the equilibrium that is a time independent solution in the given static background metric and gauge field. This equilibrium solution for the velocity field, temperature or chemical potential in terms of the background metric and gauge field can also be expanded in terms of derivatives. We shall use the following notation.

$$\begin{aligned} u^\mu|_{eq} &= \bar{u}^\mu + \delta u_{(1)}^\mu + \delta u_{(2)}^\mu + \cdots, \\ T|_{eq} &= \bar{T} + \delta T_{(1)} + \delta T_{(2)} + \cdots, \\ \nu|_{eq} &= \bar{\nu} + \delta \nu_{(1)} + \delta \nu_{(2)} + \cdots, \end{aligned} \tag{2.4}$$

where $\delta u_{(i)}^\mu$, $\delta T_{(i)}$ and $\delta \nu_{(i)}$ are i th corrections to the zeroth order equilibrium solution containing i derivatives on the background data. Now we have to substitute (2.4) in fluid stress tensor and current given in (1.4) and extract out the part that will be parity odd and which involve exactly two derivatives on the background data. This is the part which have to be equated with (2.3).

From the analysis done in [13] we know that

$$\begin{aligned}
\bar{u}^\mu &= e^{-\sigma} \{1, 0, 0, 0\}, \quad \bar{T} = T_0 e^{-\sigma}, \quad \bar{\nu} = \frac{A_0}{T_0}, \\
[\delta u_{(1)}]_0 &= 0, \quad \delta T_{(1)} = 0, \quad \delta \nu_{(1)} = 0, \\
[\delta u_{(1)}]^0 &= -a_i [\delta u_{(1)}]^i, \\
[\delta u_{(1)}]^i &= \left(\frac{b_1}{2} \right) \bar{l}^i + b_2 \bar{B}^i, \\
[\delta u_{(1)}]_i &= g_{ij} [\delta u_{(1)}]^j,
\end{aligned} \tag{2.5}$$

where

$$\begin{aligned}
F_{jk} &\equiv \partial_j A_k - \partial_k A_j, \\
\bar{l}^i &= -\frac{e^\sigma}{2} \epsilon^{ijk} f_{jk}, \\
\bar{B}^i &= \frac{1}{2} (\epsilon^{ijk} F_{jk} + A_0 \epsilon^{ijk} f_{jk}) = \frac{1}{2} \epsilon^{ijk} F_{jk} - \bar{T} \bar{\nu} \bar{l}^i, \\
b_1 &= \frac{T^3}{E + P} \left(\frac{2C\nu^3}{3} - 4C_2\nu \right), \\
b_2 &= \frac{T^2}{E + P} \left(\frac{C\nu^2}{2} - C_2 \right).
\end{aligned} \tag{2.6}$$

Here C is the anomaly coefficient and C_2 is related to the mixed anomaly in (1.3) by

$$C_2 = 8\pi^2 c_m. \tag{2.7}$$

The presence of the mixed anomaly at first order in the equilibrium partition function was pointed out in [15].

If we also expand E , P and q in terms of a derivative expansion as

$$E|_{eq} = \bar{E} + \delta E_{(1)} + \delta E_{(2)} + \dots, \quad P|_{eq} = \bar{P} + \delta P_{(1)} + \delta P_{(2)} + \dots, \quad q|_{eq} = \bar{q} + \delta q_{(1)} + \delta q_{(2)} + \dots,$$

then from (2.5) it follows that

$$\delta E_{(1)} = \delta P_{(1)} = \delta q_{(1)} = 0. \tag{2.8}$$

Using the fact that $(u_\mu u^\mu = -1)$ to all order in derivative expansion we find that

$$[\delta u_{(2)}]_0 \propto [\delta u_{(1)}]^i [\delta u_{(1)}]_i = \text{Parity even} \sim 0 \quad (\text{For our purpose}). \tag{2.9}$$

Also using the Landau gauge condition on the second order stress tensor and current one can show that on equilibrium

$$[T_{(2)}]_{00} = [T_{(2)}]_0^i = [J_{(2)}]_0 = 0. \tag{2.10}$$

Using (2.5), (2.8), (2.9) and (2.10) we get the following for the second order stress tensor and current evaluated on equilibrium.

$$\begin{aligned} [\Pi^{(2)}]_{00} &= \delta E_{(2)} \bar{u}_0^2 + \delta(-2\eta\sigma_{00}) + \delta(\zeta\Theta P_{00}), \\ [\Pi^{(2)}]_0^i &= (\bar{E} + \bar{P}) \bar{u}_0 [\delta u_{(2)}]^i + \delta(-2\eta\sigma_0^i) + \delta(\zeta\Theta P_0^i), \\ [j^{(2)}]_0 &= \delta q_{(2)} \bar{u}_0 + \delta(\Delta V_0) + \delta(\xi_l l_0) + \delta(\xi_B B_0), \end{aligned} \quad (2.11)$$

where $\delta(-2\eta\sigma_{\mu\nu})$, $\delta(\zeta\Theta P_{\mu\nu})$ and $\delta(\Delta V_\mu)$ are the two derivative corrections of $(-2\eta\sigma_{\mu\nu})$, $(\zeta\Theta P_{\mu\nu})$ and (ΔV_μ) when evaluated on $\delta u_{(1)}^\mu$, $\delta T_{(1)}$ and $\delta\nu_{(1)}$.

Now if $Q_{\mu\nu}^{(1)}$ is a tensor which is first order in the derivative expansion satisfying the following two conditions

$$Q_{\mu\nu}^{(1)} u^\mu = 0 \quad \text{at all orders and} \quad Q_{\mu\nu}^{(1)}|_{(\bar{u}^\mu, \bar{T}, \bar{\nu})} = 0, \quad (2.12)$$

then one can show in general that ⁴

$$\delta Q_{00}^{(1)} = \delta[Q^{(1)}]_0^i = 0. \quad (2.13)$$

Similarly if $Q_\mu^{(1)}$ is a vector which is first order in the derivative expansion satisfying

$$Q_\mu^{(1)} u^\mu = 0 \quad \text{at all orders and} \quad Q_\mu^{(1)}|_{(\bar{u}^\mu, \bar{T}, \bar{\nu})} = 0, \quad (2.14)$$

then it follows that ⁵

$$\delta Q_0^{(1)} = 0. \quad (2.15)$$

This argument allows us to conclude that

$$\delta(-2\eta\sigma_{00}) = \delta(\zeta\Theta P_{00}) = \delta(-2\eta\sigma_0^i) = \delta(\zeta\Theta P_0^i) = \delta(\Delta V_0) = 0.$$

Similarly $\delta(\xi_l l_\mu)$ and $\delta(\xi_B B_\mu)$ are the two derivative corrections of $(\xi_l l_\mu)$ and $(\xi_B B_\mu)$ when evaluated on $\delta u_{(1)}^\mu$, $\delta T_{(1)}$ and $\delta\nu_{(1)}$. But l_μ and B_μ are already parity odd. Therefore $\delta(\xi_l l_\mu)$ and $\delta(\xi_B B_\mu)$ are going to be parity even and hence we can set them to zero for our purpose. Putting all this together we have the following result for the components of the stress tensor and the current at the second order in derivative expansion

$$\begin{aligned} [\Pi^{(2)}]_{00} &= \delta E_{(2)} \bar{u}_0^2 = e^{2\sigma} \delta E_{(2)}, \\ [\Pi^{(2)}]_0^i &= (\bar{E} + \bar{P}) \bar{u}_0 [\delta u_{(2)}]^i = -e^\sigma (\bar{E} + \bar{P}) [\delta u_{(2)}]^i, \\ [j^{(2)}]_0 &= -e^\sigma \delta q_{(2)}. \end{aligned} \quad (2.16)$$

⁴ $0 = \delta[u^\mu Q_{\mu\nu}^{(1)}] = \bar{u}^\mu \delta Q_{\mu\nu}^{(1)} + \delta u^\mu [Q_{\mu\nu}^{(1)}|_{(\bar{u}^\alpha, \bar{T}, \bar{\nu})}] = e^{-\sigma} \delta Q_{0\nu}^{(1)}.$

⁵ $0 = \delta[u^\mu Q_\mu^{(1)}] = \bar{u}^\mu \delta Q_\mu^{(1)} + \delta u^\mu [Q_\mu^{(1)}|_{(\bar{u}^\alpha, \bar{T}, \bar{\nu})}] = e^{-\sigma} \delta Q_0^{(1)}.$

Inverting (2.16) and using (2.3) we can determine the second order piece of the equilibrium solution for velocity, temperature and chemical potential in terms of background data. Now the components $[\Pi^{(2)}]^{ij}$ and $[j^{(2)}]^i$ will give rise to the constraints on the transport coefficients.

$$\begin{aligned} [\Pi^{(2)}]^{ij} &= \delta P_2 g^{ij} - 2\eta \delta\sigma^{ij} - \zeta \delta\Theta g^{ij} + [T_{(2)}]^{ij}, \\ [j^{(2)}]^i &= \bar{q}[\delta u_{(2)}]^i + \Delta \delta[V^i] + [J_{(2)}]^i. \end{aligned} \quad (2.17)$$

We have to use (2.16) to determine δP_2 and $[\delta u_{(2)}]^i$.

The transport coefficients are determined by demanding that (2.17) is satisfied. For convenience let us further split the first equation in (2.17) in two parts, the trace part which is obtained by contracting the equation with g_{ij} and a traceless part which is obtained by subtracting the trace part of the equation from the first equation of (2.17) ⁶. These are given by

Trace part :

$$\delta P_2 - \zeta \delta\Theta + \frac{1}{3}(-2\eta \delta\sigma^{ij} + [T_{(2)}]^{ij})g_{ij} = 0. \quad (2.18)$$

Traceless part :

$$-2\eta \left(\delta\sigma^{ij} - \frac{g^{ij}}{3}(\delta\sigma^{lm}g_{lm}) \right) + \left([T_{(2)}]^{ij} - \frac{g^{ij}}{3}([T_{(2)}]^{lm}g_{lm}) \right) = 0.$$

In (2.18) we have used (2.3) to set $[\Pi^{(2)}]^{ij}$ to zero. Therefore the RHS of (2.18) vanishes.

Transport in the traceless part of the stress tensor

In this subsection we shall analyze the second equation of (2.18).

Traceless part :

$$-2\eta \left(\delta\sigma^{ij} - \frac{g^{ij}}{3}(\delta\sigma^{lm}g_{lm}) \right) + \left([T_{(2)}]^{ij} - \frac{g^{ij}}{3}([T_{(2)}]^{lm}g_{lm}) \right) = 0. \quad (2.19)$$

We now go through the following steps:

- We first evaluate the second order contribution from σ^{ij} .
- We then write down the most general form for $[T_{(2)}]^{(\mu\nu)}$ from usual symmetry analysis and on shell independence. In general it will contain many unknown transport coefficients to begin with. But if we restrict our attention to only the parity odd ones, the total number of terms is 18 (Φ_i , ($i = 1, \dots, 12$) multiplying 12 independent traceless symmetric tensors and χ , ($i = 1, \dots, 6$) multiplying 6 independent scalars, see (1.8)). All these 18 terms have been listed in table 2.

⁶Note that here trace is taken with the 3 dimensional metric.

- One can see that all the 6 χ_i s are not going to appear in the combination that we are going to evaluate in (2.19).
- So we have to evaluate the rest of the 12 terms (multiplying Φ_i s on the equilibrium time-independent solution).

For this it is sufficient to substitute only the zeroth order equilibrium solution, since each of these terms already contain two derivatives.

- As mentioned in section 1.2, it turns out that 6 of these twelve terms evaluate to zero on equilibrium. So there are only 6 transport coefficients which can be constrained by this equilibrium analysis. These 6 terms are the following.

$$\begin{aligned}\tau_{\mu\nu}^{(1)} &= \bar{\nabla}_{\langle\mu} l_{\nu\rangle}, \quad \tau_{\mu\nu}^{(2)} = \bar{\nabla}_{\langle\mu} B_{\nu\rangle}, \quad \tau_{\mu\nu}^{(3)} = (\partial_{\langle\mu} \nu) l_{\nu\rangle}, \\ \tau_{\mu\nu}^{(4)} &= (\partial_{\langle\mu} \nu) B_{\nu\rangle}, \quad \tau_{\mu\nu}^{(5)} = (\partial_{\langle\mu} T) l_{\nu\rangle}, \quad \tau_{\mu\nu}^{(6)} = (\partial_{\langle\mu} T) B_{\nu\rangle},\end{aligned}\tag{2.20}$$

where $l^\mu \equiv \epsilon^{\mu\nu\alpha\beta} u_\nu \partial_\alpha u_\beta$ and $B^\mu \equiv \epsilon^{\mu\nu\alpha\beta} u_\nu \partial_\alpha A_\beta$. Let us now rewrite the traceless part of the second order fluid stress tensor with these six terms, this is given by

$$[T_{(2)}]^{\mu\nu}_{odd} = \sum_{a=1}^6 \Phi_a [\tau^{(a)}]^{\mu\nu} + \text{trace part} + \text{terms that vanish on equilibrium}.\tag{2.21}$$

- From (2.19) we see that each of these six Φ_a 's has to be such that, they cancel the contribution of $\delta\sigma_{\mu\nu}$ when all of them are evaluated on equilibrium.
- Evaluating the spatial components of these six terms on the zeroth order equilibrium solution given in (2.5) we obtain

$$\begin{aligned}[\tau^{(1)}]^{ij} &= \bar{\nabla}^{(i} l^{j)} = g^{il} g^{jm} \left[\frac{\nabla_l \bar{l}_m + \nabla_m \bar{l}_l}{2} - \frac{g_{lm}}{3} (\nabla_k \bar{l}^k) - \frac{g_{lm}}{3} (\partial_k \sigma) \bar{l}^k \right] + \mathcal{O}(\partial^3), \\ [\tau^{(2)}]^{ij} &= \bar{\nabla}^{(i} B^{j)} = g^{il} g^{jm} \left[\frac{\nabla_l \bar{B}_m + \nabla_m \bar{B}_l}{2} - \frac{g_{lm}}{3} (\nabla_k \bar{B}^k) - \frac{g_{lm}}{3} (\partial_k \sigma) \bar{B}^k \right] + \mathcal{O}(\partial^3), \\ [\tau^{(3)}]^{ij} &= (\bar{\nabla}^{\langle\mu} \nu) l^{\nu\rangle} = g^{il} g^{jm} \left[\frac{(\nabla_l \bar{\nu}) \bar{l}_m + (\nabla_m \bar{\nu}) \bar{l}_l}{2} - \frac{g_{lm}}{3} (\nabla_k \bar{\nu} \bar{l}^k) \right] + \mathcal{O}(\partial^3), \\ [\tau^{(4)}]^{ij} &= (\bar{\nabla}^{\langle\mu} \nu) B^{\nu\rangle} = g^{il} g^{jm} \left[\frac{(\nabla_l \bar{\nu}) \bar{B}_m + (\nabla_m \bar{\nu}) \bar{B}_l}{2} - \frac{g_{lm}}{3} (\bar{B}^k \nabla_k \bar{\nu}) \right] + \mathcal{O}(\partial^3), \\ [\tau^{(5)}]^{ij} &= (\bar{\nabla}^{\langle\mu} T) l^{\nu\rangle} = g^{il} g^{jm} \left[\frac{(\nabla_l \bar{T}) \bar{l}_m + (\nabla_m \bar{T}) \bar{l}_l}{2} - \frac{g_{lm}}{3} (\bar{l}^k \nabla_k \bar{T}) \right] + \mathcal{O}(\partial^3), \\ [\tau^{(6)}]^{ij} &= (\bar{\nabla}^{\langle\mu} T) B^{\nu\rangle} = g^{il} g^{jm} \left[\frac{(\nabla_l \bar{T}) \bar{B}_m + (\nabla_m \bar{T}) \bar{B}_l}{2} - \frac{g_{lm}}{3} (\bar{B}^k \nabla_k \bar{T}) \right] + \mathcal{O}(\partial^3),\end{aligned}\tag{2.22}$$

where $\bar{l}^i \equiv -\frac{e^\sigma}{2} \epsilon^{ijk} f_{jk}$ and $\bar{B}^i \equiv \frac{1}{2} \epsilon^{ijk} (F_{jk} + A_0 f_{jk})$. To evaluate equation (2.22) we have extensively used section 2 of [13].

- The spatial components of the shear tensor gives the following contribution to $[\Pi^{(2)}]^{ij}$

$$\begin{aligned}
& -2\eta \delta\sigma^{ij} \\
& = -2\eta \left[\bar{P}^{i\mu} \bar{P}^{j\nu} \left\{ \frac{\bar{\nabla}_\mu [\delta u_{(1)}]_\nu + \bar{\nabla}_\nu [\delta u_{(1)}]_\mu}{2} - \frac{\bar{\nabla}_\alpha [\delta u_{(1)}]^\alpha}{3} \right\} \right. \\
& \quad \left. + (\bar{P}^{j\mu} \delta P^{i\nu} + \bar{P}^{i\nu} \delta P^{j\mu}) \left(\frac{\bar{\nabla}_\mu \bar{u}_\nu + \bar{\nabla}_\nu \bar{u}_\mu}{2} \right) \right] \\
& = -2\eta g^{il} g^{jm} \left[\left[\frac{\nabla_l [\delta u_{(1)}]_m + \nabla_m [\delta u_{(1)}]_l}{2} - \frac{g_{lm}}{3} \nabla_k [\delta u_{(1)}]^k - \frac{g_{lm}}{3} (\partial_k \sigma) [\delta u_{(1)}]^k \right. \right. \\
& \quad \left. \left. + \left[\frac{\delta u^i g^{jk} (\partial_k \sigma) + [\delta u_{(1)}]^j g^{ik} (\partial_k \sigma)}{2} \right] \right] \right], \tag{2.23}
\end{aligned}$$

where

$$\bar{P}^{i\mu} = \bar{u}^i \bar{u}^\mu + G^{i\mu} = G^{i\mu}, \quad \bar{P}^{j\mu} \delta P^{i\nu} = G^{j\mu} \bar{u}^\nu [\delta u_{(1)}]^i.$$

The last term in the last line of (2.23) results from this expansion of the projectors. Substituting (2.5) in (2.23) we obtain the following

$$\begin{aligned}
-2\eta \delta\sigma^{ij} & = -2\eta \left[\frac{b_1}{2} [\tau^{(1)}]^{ij} + b_2 [\tau^{(2)}]^{ij} + \frac{1}{2} \left(\frac{\partial b_1}{\partial \nu} \right) [\tau^{(3)}]^{ij} + \left(\frac{\partial b_2}{\partial \nu} \right) [\tau^{(4)}]^{ij} \right. \\
& \quad \left. + \frac{1}{2} \left(-\frac{b_1}{T} + \frac{\partial b_1}{\partial T} \right) [\tau^{(5)}]^{ij} + \left(-\frac{b_2}{T} + \frac{\partial b_2}{\partial T} \right) [\tau^{(6)}]^{ij} \right]. \tag{2.24}
\end{aligned}$$

Next we have to substitute (2.23) and (2.24) in (2.19). Now to satisfy (2.19) the coefficient of each independent expression should vanish. From examining (2.23) and (2.24) it seems that demanding every independent term to vanish results in more equations than the unknowns which are the transport coefficients. However from the structure of the equations in (2.19) it is clear that the equation admits the following unique solution for the transport coefficients

$$\begin{aligned}
\Phi_1 & = \eta b_1, \quad \Phi_2 = 2\eta b_2, \quad \Phi_3 = \eta \left(\frac{\partial b_1}{\partial \nu} \right), \quad \Phi_4 = 2\eta \left(\frac{\partial b_2}{\partial \nu} \right), \\
\Phi_5 & = \eta \left[-\frac{b_1}{T} + \frac{\partial b_1}{\partial T} \right], \quad \Phi_6 = 2\eta \left[-\frac{b_2}{T} + \frac{\partial b_2}{\partial T} \right], \tag{2.25}
\end{aligned}$$

where

$$b_1 = \frac{T^3}{E+P} \left(\frac{2C\nu^3}{3} - 4C_2\nu \right), \quad b_2 = \frac{T^2}{E+P} \left(\frac{C\nu^2}{2} - C_2 \right), \tag{2.26}$$

and E, P, T is the energy density, pressure and temperature respectively. $\nu = \frac{\mu}{T}$ refers to the chemical potential, and η the shear viscosity. C is the gauge anomaly coefficient, while C_2 is related to the mixed anomaly by (1.11). The transport coefficient Φ_1 gives rise to chiral dispersion relations in the shear mode [12].

Transport in the trace of the stress tensor and current

We will now constrain the transport coefficients that occur in the trace part of the stress tensor and the current using the equilibrium partition function. We will first rewrite relevant part of the current and the trace part of the stress tensor up to 2nd order in derivative expansion. Using (2.3) and (2.17) we obtain the following equation for the trace part of the stress tensor and current at second order.

$$\begin{aligned}
\text{Trace part : } 0 &= \delta P_2 - \zeta \delta \Theta + \frac{1}{3}(-2\eta \delta \sigma^{ij} + [T_{(2)}]^{ij})g_{ij} \\
&= \delta P_2 - \zeta \delta \Theta + \left[\sum_1^6 \chi_i \mathcal{S}_i \right]_{eq} \\
&= \left[-\zeta \delta \Theta + \left(\frac{\partial P}{\partial E} \right)_q \delta E_{(2)} + \left(\frac{\partial P}{\partial q} \right)_E \delta q_{(2)} \right] + \left[\sum_1^6 \chi_i \mathcal{S}_i \right]_{eq}
\end{aligned} \tag{2.27}$$

$$\begin{aligned}
\text{Current : } [J_{(2)}]^i &= [j^{(2)}]^i - \delta[\Delta V^i] - q[\delta u_{(2)}]^i \\
\Rightarrow \left[\sum_{k=1}^9 \delta_k \mathcal{V}_{(k)}^i \right]_{eq} &= \bar{T} \left(\frac{\partial M_1}{\partial \bar{T}} \right) \epsilon^{ijk} (\partial_j \bar{T}) (\partial_k \bar{\nu}) - \delta[\Delta V^i] - q[\delta u_{(2)}]^i.
\end{aligned} \tag{2.28}$$

Both in (2.27) and (2.28) we have used (1.8) to write the fluid stress tensor and the current in terms of the transport coefficients. Also in (2.27) to evaluate the combination $(-2\eta \delta \sigma^{ij} + [T_{(2)}]^{ij})g_{ij}$ we have used (2.24) and (2.25). One can see that the transport coefficients Φ_i s drop out.

Using (2.16) and (2.3) we evaluate δE_2 , δq_2 and $[\delta u_{(2)}]^i$, this results in

$$\begin{aligned}
\delta E_2 &= e^{-2\sigma} \Pi_{00}^{(2)} = 2\bar{T}^2 \left(\frac{\partial M_1}{\partial \bar{T}} \right) (\bar{B}^i \partial_i \bar{\nu}) - 2T^3 \left(\frac{\partial M_2}{\partial \bar{T}} - \bar{\nu} \frac{\partial M_1}{\partial \bar{T}} \right) (\bar{l}^i \partial_i \bar{\nu}), \\
\delta q_2 &= e^{-\sigma} j_0^{(2)} 2 \left(\frac{\partial M_1}{\partial \bar{T}} \right) (\bar{B}^i \partial_i \bar{T}) - 2\bar{T} \left(\frac{\partial M_2}{\partial \bar{T}} - \bar{\nu} \frac{\partial M_1}{\partial \bar{T}} \right) (\bar{l}^i \partial_i \bar{T}), \\
[\delta u_{(2)}]^i &= - \left(\frac{e^{-\sigma}}{\bar{E} + \bar{P}} \right) [\Pi^{(2)}]_0^i = - \left(\frac{2\bar{T}^2}{\bar{E} + \bar{P}} \right) \left(\frac{\partial M_2}{\partial \bar{T}} - \bar{\nu} \frac{\partial M_1}{\partial \bar{T}} \right) \epsilon^{ijk} (\partial_j \bar{T}) (\partial_k \bar{\nu}).
\end{aligned} \tag{2.29}$$

Now we have to compute $\delta \Theta$ and $\delta(\Delta V^i)$.

$$\begin{aligned}
\delta \Theta &= \bar{\nabla}_\mu [\delta u_{(1)}]^\mu = \nabla_k [\delta u_{(1)}]^k - \left(\frac{\partial_k \bar{T}}{\bar{T}} \right) [\delta u_{(1)}]^k \\
&= \frac{1}{2} \left(\frac{\partial b_1}{\partial \bar{T}} - \frac{2b_1}{\bar{T}} \right) (\bar{l}^i \partial_i \bar{T}) + \frac{1}{2} \left(\frac{\partial b_1}{\partial \bar{\nu}} - 2b_2 \bar{T} \right) (\bar{l}^i \partial_i \bar{\nu}) \\
&\quad + \left(\frac{\partial b_2}{\partial \bar{T}} - \frac{b_2}{\bar{T}} \right) (\bar{B}^i \partial_i \bar{T}) + \left(\frac{\partial b_2}{\partial \bar{\nu}} \right) (\bar{B}^i \partial_i \bar{\nu}).
\end{aligned} \tag{2.30}$$

To derive (2.30) we have used the following identities

$$\nabla_i \bar{l}^i = - \left(\frac{\partial_i \bar{T}}{\bar{T}} \right) \bar{l}^i, \quad \nabla_i \bar{B}^i = - \bar{T} \bar{l}^i \partial_i \bar{\nu}. \quad (2.31)$$

Similarly δV^i is given by the following expression

$$\delta \{ \Delta V^i \} = \Delta G^{i\mu} \mathcal{F}_{\mu\nu} \delta u^\nu = \Delta \epsilon^{ijk} [\delta u_{(1)}]_j \bar{B}_k = \Delta \left(\frac{b_1}{2} \right) \epsilon^{ijk} \bar{l}_j \bar{B}_k, \quad (2.32)$$

where $\mathcal{F}_{\mu\nu} = \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu$.

Now we have to evaluate the 6 scalars ($\mathcal{S}_i$, $i = 1, \dots, 6$) and the i th component of the 9 vectors ($\mathcal{V}_{(k)}^i$, $k = 1, \dots, 9$) on the equilibrium solution. We need the equilibrium solution only at zeroth order since all of them already contain two derivatives. Explicit expressions for all these terms are listed in table 2.

As mentioned in section 1.2, only 6 terms, 4 scalars and 2 vectors are non-vanishing on the zeroth order equilibrium solution. Thus the relevant parts of the second order current and trace of the stress tensor are given by

$$\begin{aligned} [T_{(2)}]^{\alpha\beta}|_{trace \ part} &= P^{\alpha\beta} [\chi_1 (l^\mu \partial_\mu \nu) + \chi_2 (B^\mu \partial_\mu \nu) + \chi_3 (l^\mu \partial_\mu T) + \chi_4 (B^\mu \partial_\mu T)], \\ J_{(2)}^\mu &= \Delta_1 [\epsilon^{\mu\nu\alpha\beta} u_\nu (\partial_\alpha \nu) (\partial_\alpha T)] + \Delta_2 [\epsilon^{\mu\nu\alpha\beta} u_\nu B_\alpha l_\beta]. \end{aligned} \quad (2.33)$$

We then evaluate all these six terms on the zeroth order equilibrium solution.

$$\begin{aligned} (l^\mu \partial_\mu \nu) &= (\bar{l}^i \partial_i \bar{\nu}), \\ (B^\mu \partial_\mu \nu) &= (\bar{B}^i \partial_i \bar{\nu}), \\ (l^\mu \partial_\mu T) &= (\bar{l}^i \partial_i \bar{T}), \\ (B^\mu \partial_\mu T) &= (\bar{B}^i \partial_i \bar{T}), \\ \epsilon^{i\nu\alpha\beta} u_\nu (\partial_\alpha \nu) (\partial_\alpha T) &= \epsilon^{ijk} (\partial_j \bar{\nu}) (\partial_k \bar{T}), \\ \epsilon^{i\nu\alpha\beta} u_\nu B_\alpha l_\beta &= \epsilon^{ijk} \bar{B}_j \bar{l}_k. \end{aligned} \quad (2.34)$$

Substituting (2.34) in (2.33) we express the LHS of (2.33) in terms of the background data and the two arbitrary functions M_1 and M_2 of the second order parity odd partition function. Now using (2.27) and (2.3) we can express the 5 transport coefficients appearing in (2.33) in terms of M_1 and M_2 . Since one cannot generate a term of the form $\epsilon^{ijk} l_j B_k$ from the partition function, the 6th transport coefficients Δ_2 will be completely determined by the correction of the first order current. This is similar to the way the transport coefficients Φ_i 's, which appear in the traceless part

of the stress tensor were determined. The end result of this step is

$$\begin{aligned}
\Delta_2 &= -\frac{\Delta b_1}{2}, \\
\chi_1 &= -2R_1 T^3 \left[\bar{\nu} \frac{\partial M_1}{\partial \bar{T}} - \frac{\partial M_2}{\partial \bar{T}} \right] + \zeta \left(\frac{1}{2} \frac{\partial b_1}{\partial \nu} - b_2 \bar{T} \right), \\
\chi_2 &= -2R_1 T^2 \left(\frac{\partial M_1}{\partial \bar{T}} \right) + \zeta \left(\frac{\partial b_2}{\partial \nu} \right), \\
\chi_3 &= -2R_2 T \left[\bar{\nu} \frac{\partial M_1}{\partial T} - \frac{\partial M_2}{\partial \bar{T}} \right] + \frac{\zeta}{2} \left(\frac{\partial b_1}{\partial \bar{T}} - \frac{2b_1}{\bar{T}} \right), \\
\chi_4 &= 2R_2 \left(\frac{\partial M_1}{\partial \bar{T}} \right) + \zeta \left(\frac{\partial b_2}{\partial T} - \frac{b_2}{T} \right), \\
\Delta_1 &= 2\bar{T} \left(\frac{\partial M_1}{\partial \bar{T}} \right) + \left(\frac{2qT^2}{E+P} \right) \left(\frac{\partial M_2}{\partial \bar{T}} - \bar{\nu} \frac{\partial M_1}{\partial \bar{T}} \right),
\end{aligned} \tag{2.35}$$

where

$$\begin{aligned}
R_1 &= \left(\frac{\partial P}{\partial E} \right)_q, \quad R_2 = \left(\frac{\partial P}{\partial q} \right)_E, \\
b_1 &= \frac{T^3}{E+P} \left(\frac{2C\nu^3}{3} - 4C_2\nu \right), \quad b_2 = \frac{T^2}{E+P} \left(\frac{C\nu^2}{2} - C_2 \right).
\end{aligned}$$

Eliminating M_1 and M_2 from these expressions we get three relations among the remaining 5 second order transport coefficients.

$$\begin{aligned}
\Delta_2 &= -\frac{\Delta b_1}{2}, \\
T^2 R_1 \left[\chi_3 - \frac{\zeta}{2} \left(\frac{\partial b_1}{\partial T} - \frac{2b_1}{T} \right) \right] - R_2 \left[\chi_1 - \frac{\zeta}{2} \left(\frac{\partial b_1}{\partial \nu} - 2b_2 T \right) \right] &= 0, \\
T^2 R_1 \left[\chi_4 - \zeta \left(\frac{\partial b_2}{\partial T} - \frac{b_2}{T} \right) \right] + R_2 \left[\chi_2 - \zeta \left(\frac{\partial b_1}{\partial \nu} \right) \right] &= 0, \\
R_1 T \Delta_1 + \left[\chi_2 - \zeta \left(\frac{\partial b_2}{\partial \nu} \right) \right] - \frac{q}{(E+P)} \left[\chi_1 - \frac{\zeta}{2} \left(\frac{\partial b_1}{\partial \nu} - 2b_2 T \right) \right] &= 0.
\end{aligned} \tag{2.36}$$

3. Kubo formula for the transport coefficients Φ_1, Φ_2

In this section we derive the relations obtained for the transport coefficients Φ_1 and Φ_2 given in (2.25) using the Kubo formula. We consider the following equilibrium background for the metric, gauge field and the velocity

$$g_{\mu\nu}^{(0)} = \text{diag}(-1, 1, 1, 1), \quad \mathcal{A}_\mu = (\nu^{(0)} T_0, 0, 0, 0), \quad u^\mu = (1, 0, 0, 0). \tag{3.1}$$

The chemical potential $\nu^{(0)}$ and the temperature T_0 are constants and do not depend on space-time. Since the energy $E^{(0)}$ and the pressure $P^{(0)}$ can be thought of as functions of the temperature and the chemical potential, they are also constants in

space-time. Now consider the following non-zero metric perturbations about this background

$$\delta g_{tx} = h_{tx}, \quad \delta g_{tz} = h_{tz}, \quad \delta g_{yx} = h_{yx}, \quad \delta g_{yz} = h_{yz}. \quad (3.2)$$

The gauge field perturbations are given by

$$\delta \mathcal{A}_\mu = (0, a_x, 0, a_z). \quad (3.3)$$

The fluid velocity is close to the rest frame and its perturbations are given by

$$\delta u_\mu = (0, v^x, 0, v^z). \quad (3.4)$$

All perturbations are assumed to have dependence only in the time t and y -direction.

To derive Kubo formulae for transport coefficients, we consider the constitutive relations for the stress tensor and the charge current as one point functions in the presence of external sources. We then obtain two point functions for the currents by differentiating with respect to the metric and the gauge field perturbations. Working this out to the linear order in perturbations will result in Kubo formulae for the transport coefficients Φ_1, Φ_2 .

To proceed we will require the the Christoffel symbols to the linear order in perturbations. The non-vanishing elements at the linear order are given by

$$\begin{aligned} \Gamma_{xy}^t &= -\frac{1}{2}(\partial_y h_{tx} - \partial_t h_{yx}), & \Gamma_{zy}^t &= -\frac{1}{2}(\partial_y h_{tz} - \partial_t h_{zy}), \\ \Gamma_{tt}^x &= \partial_t h_{tx}, & \Gamma_{ty}^x &= \frac{1}{2}(\partial_y h_{tx} + \partial_t h_{yx}), & \Gamma_{yy}^x &= \partial_y h_{xy}, \\ \Gamma_{tt}^z &= \partial_t h_{tz}, & \Gamma_{ty}^z &= \frac{1}{2}(\partial_y h_{tz} + \partial_t h_{zy}), & \Gamma_{yy}^z &= \partial_y h_{yz}, \\ \Gamma_{tx}^y &= \frac{1}{2}(\partial_t h_{yx} - \partial_y h_{tx}), & \Gamma_{tz}^y &= \frac{1}{2}(\partial_t h_{yz} - \partial_y h_{tz}). \end{aligned} \quad (3.5)$$

Evaluating the inverse metric to the linear order we obtain

$$h^{tx} = h_{tx}, \quad h^{tz} = h_{tz}, \quad h^{yx} = -h_{yx}, \quad h^{yz} = -h_{yz}. \quad (3.6)$$

The covariant components of the velocity are given by

$$u_\mu = (-1, v^x + h_{xt}, 0, v^z + h_{zt}). \quad (3.7)$$

We now evaluate various components of the stress tensor to the linear order in the perturbations. From the list of terms that contribute at 2nd order in derivatives given in table 2 we see that in the background we have chosen all the scalars $\mathcal{S}_i$ vanish. The reason for this is for the background all the thermodynamic functions are independent of space-time. We now examine the traceless part of the stress tensor. Note that the contributions from $\tau_{\mu\nu}^{(i)}$ for $i = 3, 4, 5, 6, 7, 8, 9, 10, 11$ vanish since the thermodynamic functions are independent of space-time. What remains

to be evaluated are the contributions from $\tau_\mu^{(i)}$ for $i = 1, 2$ and $i = 12$. Let us first examine the tx and the ty component of the stress tensor. To the second order in derivatives and to the linear order in the perturbation this is given by

$$\begin{aligned} T^{tx} &= (E^{(0)} + P^{(0)})v^x + P^{(0)}h_{tx}, \\ T^{tz} &= (E^{(0)} + P^{(0)})v^z + P^{(0)}h_{tz}, \\ T^{ty} &= 0. \end{aligned} \tag{3.8}$$

Note that $\sigma^{tx}, \sigma^{tz}, [\tau^{(i)}]^{tx}$ and $[\tau^{(i)}]^{tz}$ for $i = 1, 2, 12$ do not contribute at the linear order. The reason is that the term $\nabla_\alpha u_\beta$ and $\nabla_\alpha l_\beta$ is a first order term, therefore one has to evaluate the projector for these components say $P^{t\alpha}P^{x\beta}$ at the zeroth order, which vanishes.

Lets examine the yx component of the stress tensor. We need to evaluate the contributions from $[\tau^{(i)}]^{yx}$ for $i = 1, 2, 12$. These are given by ⁷

$$\begin{aligned} [\tau^{(1)}]^{yx} &= \frac{1}{2}\partial_y^2(v^z + h_{zt}), \\ [\tau^{(2)}]^{yx} &= \frac{1}{2}\partial_y^2 a_z, \\ [\tau^{(12)}]^{yx} &= \frac{1}{2}\partial_y(\partial_y v^z + \partial_t h_{yz}). \end{aligned} \tag{3.9}$$

We also need the contribution of the shear tensor to the linear order. This is given by

$$\sigma^{yx} = \frac{1}{2}(\partial_y v^x + \partial_t h_{yx}). \tag{3.10}$$

Considering all these contributions along with the contribution to the stress tensor to the zeroth order in derivative we obtain

$$\begin{aligned} T^{yx} &= -P^{(0)}h_{yx} - \eta(\partial_y v^x + \partial_t h_{yx}) \\ &\quad + \frac{1}{2}\Phi_1\partial_y^2(v^z + h_{zt}) + \frac{1}{2}\Phi_2\partial_y^2 a_z \\ &\quad + \frac{1}{2}\partial_y(\partial_y v^z + \partial_t h_{yz}). \end{aligned} \tag{3.11}$$

The equations of motion for the x component of the stress tensor to the linear order in the fields is given by

$$\partial_t T^{tx} + \partial_t h_{tx} T^{tt} + \partial_y T^{yx} = 0. \tag{3.12}$$

Here T^{tt} is the zeroth order tt component of the stress tensor which is given by

$$T^{tt} = E^{(0)}. \tag{3.13}$$

⁷In evaluating these contributions we take $\epsilon^{01234} = \frac{1}{\sqrt{-g}}$.

Fourier transforming these equations and taking the zero frequency limit or taking the time independent situation we obtain the Ward identities for the one point function of the stress tensor.

$$\lim_{\omega \rightarrow 0} T^{yx}(k) = 0. \quad (3.14)$$

We can now differentiate these with respect to the background fields h_{zt} , and a_z and obtain Kubo formulae for the transport coefficients Φ_1 and Φ_2 respectively. A similar procedure for the yz component of the stress tensor yields the same result.

To proceed we first eliminate v^x and v^z using (3.8). This results in the following equations

$$\begin{aligned} T^{yx} = & -P^{(0)} h_{yx} - \eta \left(\frac{\partial_y T^{tx} - P^{(0)} \partial_y h_{tx}}{E^{(0)} + P^{(0)}} \right) \\ & + \frac{\Phi_1}{2} \left(\frac{\partial_y^2 T^{tz} + E^{(0)} \partial_y^2 h_{tz}}{E^{(0)} + P^{(0)}} \right) + \frac{\Phi_2}{2} \partial_y^2 a_z \\ & + \frac{\Phi_{12}}{2} \left(\frac{\partial_y^2 T^{tz} - P^{(0)} \partial_y^2 h_{tz}}{E^{(0)} + P^{(0)}} \right). \end{aligned} \quad (3.15)$$

Here we have already used time-independence to drop the time derivatives. Fourier transforming these equations we obtain

$$\begin{aligned} T^{yx}(k) = & -P^{(0)} h_{yx} - \eta \left(\frac{ik T^{tx} - ik P^{(0)} h_{tx}}{E^{(0)} + P^{(0)}} \right) \\ & - \frac{\Phi_1}{2} \left(\frac{k^2 T^{tz} + E^{(0)} k^2 h_{tz}}{E^{(0)} + P^{(0)}} \right) - \frac{\Phi_2}{2} k^2 a_z \\ & - \frac{\Phi_{12}}{2} k^2 \left(\frac{T^{tz} - P^{(0)} h_{tz}}{E^{(0)} + P^{(0)}} \right) \\ = & 0. \end{aligned} \quad (3.16)$$

The last equality in the above equation implements the Ward identity given in (3.14) and k is the momentum in the y direction. Let us now focus on the expression for T^{yx} a similar analysis can be repeated for T^{yz} . Differentiating the Ward identity for T^{yx} with respect to h_{zt} and a_z and setting the other backgrounds to zero we obtain the following two equations

$$\begin{aligned} \frac{k^2}{E^{(0)} + P^{(0)}} [(\Phi_1 \langle T^{tz}(k) T^{tz}(-k) \rangle + E^{(0)}) + \Phi_{12} (\langle T^{tz}(k) T^{tz}(-k) \rangle - P^{(0)})] \\ = -ik \frac{2\eta}{E^{(0)} + P^{(0)}} \langle T^{tx}(k) T^{tz}(-k) \rangle, \\ \frac{\Phi_1 + \Phi_{12}}{E^{(0)} + P^{(0)}} k^2 \langle T^{tz}(k) j^z(-k) \rangle + \Phi_2 k^2 = -ik \frac{2\eta}{E^{(0)} + P^{(0)}} \langle T^{tz}(k) j^z(-k) \rangle. \end{aligned} \quad (3.17)$$

To obtain the first equation we have differentiated with respect to h_{tz} and set all the other backgrounds to zero. The second equation is obtained by differentiating

the Ward identity with respect to a_z and setting the remaining backgrounds to zero. The equations in (3.17) are sufficient to determine the transport coefficients Φ_1, Φ_2 . From [8] we have the following results for the various two point functions.

$$\begin{aligned}\lim_{k \rightarrow 0, \omega \rightarrow 0} \langle T^{tx}(k) T^{tz}(-k) \rangle &= ik \left(\frac{C}{3} (\nu^{(0)} T^{(0)})^3 - 2C_2 (T^{(0)})^3 \nu^{(0)} \right), \\ \lim_{k \rightarrow 0, \omega \rightarrow 0} \langle T^{tx}(k) j^z(-k) \rangle &= ik \left(\frac{C}{2} (\nu^{(0)} T^{(0)})^2 - C_2 (T^{(0)})^2 \right).\end{aligned}\quad (3.18)$$

These results are given in equations (79)-(81) for a system with 3 chemical potentials and equations (123)-(125) for a system with a single chemical potential of [8]. The definition of the variables for the two point functions used is given in equation (47). This reference also uses the normalization

$$\frac{-C}{8} = \frac{1}{32\pi^2}, \quad \frac{c_m}{4} = \frac{1}{768\pi^2}, \quad C_2 = \frac{1}{24}, \quad (3.19)$$

for the gauge anomaly and we have rewritten the chemical potential μ in terms of the variable ν . From [16] we can read out the following correlators

$$\lim_{k \rightarrow 0, \omega \rightarrow 0} \langle T^{tz}(k) T^{tz}(-k) \rangle = P^{(0)}, \quad \lim_{k \rightarrow 0, \omega \rightarrow 0} \langle T^{tz}(k) j^z(-k) \rangle = 0. \quad (3.20)$$

These correlators are mentioned below equation (2.16) of reference [16]. Substituting the formulae for the two point functions given in (3.18) and (3.20) into the equations given in (3.17) we obtain

$$\begin{aligned}\Phi_1 &= \frac{2\eta}{E^{(0)} + P^{(0)}} \left(\frac{C}{3} (\nu^{(0)} T^{(0)})^3 - 2C_2 (T^{(0)})^3 \nu^{(0)} \right), \\ \Phi_2 &= \frac{2\eta}{E^{(0)} + P^{(0)}} \left(\frac{C}{2} (\nu^{(0)} T^{(0)})^2 - C_2 (T^{(0)})^2 \right).\end{aligned}\quad (3.21)$$

These expressions agree with that derived using the equilibrium partition function method which are given in (2.25).

4. Conclusions

We have used the equilibrium partition function to obtain expressions for 7 parity odd transport coefficients which occur at 2nd order for a non-conformal fluid with a single conserved charge. These transport coefficients can be expressed in terms of the anomaly, shear viscosity, charge diffusivity and thermodynamic functions. Out of these 2 transport coefficients can also be derived using the Kubo formulae. These formulae agree with that obtained by the partition function method. The equilibrium partition function also gives 3 constraints for 5 other parity odd transport coefficients at this order. The transport coefficient Φ_1 affects chiral dispersion relations [12].

As we have mentioned earlier, parity odd coefficients have been examined earlier for conformal charge transport in [12]. There the principle used was that these coefficients should not contribute to entropy production. In general the constraints obtained by examining the zero entropy production condition will be more than that obtained from the equilibrium partition function. It will be interesting to carry out the analysis of [12] to non-conformal fluids and compare with the results obtained in this paper.

Our analysis of these transport properties were motivated due to the possibility of studying them in holography. While this work was being done we received preprint [17] which evaluates all 2nd order transport coefficients for a charged conformal fluid in the framework of AdS/CFT. It is useful to compare the relations we have obtained for the parity odd sector with the expressions of [17].

Finally it will be useful to develop Kubo like expressions for all the parity odd transport coefficients. From the constitutive relations it seems that the remaining transport coefficients involves 3 point functions. Determining these relations will provide an alternate method to obtain the transport coefficients from holography.

Acknowledgments

We thank Aninda Sinha for organizing a stimulating workshop on “Non-perturbative gauge theories, holography and all that” which initiated this collaboration. We thank Nabamita Banarjee, Bobby Ezuthachan, Yashodhan Hatwalne, Sachin Jain, Shiraz Minwalla and Rahul Pandit for useful and stimulating discussions. S.B thanks the CHEP, IISc and the ICTS-TIFR for hospitality during the early phase of this work. The work of J.R.D is partially supported by the Ramanujan fellowship DST-SR/S2/RJN-59/2009.

References

- [1] D. T. Son and P. Surowka, *Hydrodynamics with Triangle Anomalies*, *Phys.Rev.Lett.* **103** (2009) 191601, [[arXiv:0906.5044](#)].
- [2] N. Banerjee, J. Bhattacharya, S. Bhattacharyya, S. Dutta, R. Loganayagam, *et. al.*, *Hydrodynamics from charged black branes*, *JHEP* **1101** (2011) 094, [[arXiv:0809.2596](#)].
- [3] J. Erdmenger, M. Haack, M. Kaminski, and A. Yarom, *Fluid dynamics of R-charged black holes*, *JHEP* **0901** (2009) 055, [[arXiv:0809.2488](#)].
- [4] R. Loganayagam and P. Surowka, *Anomaly/Transport in an Ideal Weyl gas*, *JHEP* **1204** (2012) 097, [[arXiv:1201.2812](#)].
- [5] K. Jensen, *Triangle Anomalies, Thermodynamics, and Hydrodynamics*, *Phys.Rev.* **D85** (2012) 125017, [[arXiv:1203.3599](#)].

- [6] N. Banerjee, S. Dutta, S. Jain, R. Loganayagam, and T. Sharma, *Constraints on Anomalous Fluid in Arbitrary Dimensions*, *JHEP* **1303** (2013) 048, [[arXiv:1206.6499](#)].
- [7] R. Loganayagam, *Anomalies and the Helicity of the Thermal State*, [arXiv:1211.3850](#).
- [8] K. Landsteiner, E. Megias, and F. Pena-Benitez, *Anomalous Transport from Kubo Formulae*, [arXiv:1207.5808](#).
- [9] P. Romatschke and D. T. Son, *Spectral sum rules for the quark-gluon plasma*, *Phys.Rev.* **D80** (2009) 065021, [[arXiv:0903.3946](#)].
- [10] A. V. Andreev, D. T. Son, and B. Spivak, *Hydrodynamics of liquids of chiral molecules and suspensions containing chiral particles*, *Phys. Rev. Lett.* **104** (May, 2010) 198301.
- [11] B. Sahoo and H.-U. Yee, *Holographic chiral shear waves from anomaly*, *Phys.Lett.* **B689** (2010) 206–212, [[arXiv:0910.5915](#)].
- [12] D. E. Kharzeev and H.-U. Yee, *Anomalies and time reversal invariance in relativistic hydrodynamics: the second order and higher dimensional formulations*, *Phys.Rev.* **D84** (2011) 045025, [[arXiv:1105.6360](#)].
- [13] N. Banerjee, J. Bhattacharya, S. Bhattacharyya, S. Jain, S. Minwalla, *et. al.*, *Constraints on Fluid Dynamics from Equilibrium Partition Functions*, *JHEP* **1209** (2012) 046, [[arXiv:1203.3544](#)].
- [14] K. Jensen, M. Kaminski, P. Kovtun, R. Meyer, A. Ritz, *et. al.*, *Towards hydrodynamics without an entropy current*, *Phys.Rev.Lett.* **109** (2012) 101601, [[arXiv:1203.3556](#)].
- [15] K. Jensen, R. Loganayagam, and A. Yarom, *Thermodynamics, gravitational anomalies and cones*, *JHEP* **1302** (2013) 088, [[arXiv:1207.5824](#)].
- [16] I. Amado, K. Landsteiner, and F. Pena-Benitez, *Anomalous transport coefficients from Kubo formulas in Holography*, *JHEP* **1105** (2011) 081, [[arXiv:1102.4577](#)].
- [17] E. Megias and F. Pena-Benitez, *Holographic Gravitational Anomaly in First and Second Order Hydrodynamics*, [arXiv:1304.5529](#).